

1233 and 1224 respectively) were used (50 pmoles each) with the pGEM3z(f+) plasmid vector (Promega Corp.) as template (10 ng) containing the target sequences. The conditions for PCR were as follows: 50 μ M of each dNTP and 2.5 units of Taq DNA polymerase in 100 μ l of 20 mM Tris-Cl, pH 8.3, 1.5 mM MgCl₂, 50 mM KCl with 0.05% Tween-20 and 0.05% NP-40. Reactions were cycled 35 times through 95°C for 45 seconds, 63°C for 45 seconds, then 72°C for 75 seconds. After cycling, reactions were finished off with an incubation at 72°C for 5 minutes. The resulting fragment was purified by electrophoresis through a 6% polyacrylamide gel (29:1 cross link) in a buffer of 45 mM Tris-Borate, pH 8.3, 1.4 mM EDTA, visualized by ethidium bromide staining or autoradiography, excised from the gel, eluted by passive diffusion, and concentrated by ethanol precipitation.

Cleavage of the substrate DNA occurred in the presence of the pilot oligonucleotide 19-12 at 50°C (Fig. 12B, lanes 1 and 7) but not at 75°C (lanes 4 and 10). In the presence of oligonucleotide 30-12 cleavage was observed at both temperatures. Cleavage did not occur in the absence of added oligonucleotides (lanes 3, 6 and 12) or at about 80°C even though at 50°C adventitious structures in the substrate allowed primer-independent cleavage in the absence of KCl (Fig. 12B, lane 9). A non-specific oligonucleotide with no complementarity to the substrate DNA did not direct cleavage at 50°C, either in the absence or presence of 50 mM KCl (lanes 13 and 14). Thus, the specificity of the cleavage reactions can be controlled by the extent of complementarity to the substrate and by the conditions of incubation.

D. Cleavage Of RNA

An shortened RNA version of the sequence used in the transcleavage experiments discussed above was tested for its ability to serve as a substrate in the reaction. The RNA is cleaved at the expected place, in a reaction that is dependent upon the presence of the pilot oligonucleotide. The RNA substrate, made by T7 RNA polymerase in the presence of [α -³²P]UTP, corresponds to a truncated version of the DNA substrate used in Figure 12B. Reaction conditions were similar to those in used for the DNA substrates described above, with 50 mM KCl; incubation was for 40

minutes at 55°C. The pilot oligonucleotide used is termed 30-0 (SEQ ID NO:20) and is shown in Fig. 13A.

The results of the cleavage reaction is shown in Figure 13B. The reaction was run either in the presence or absence of DNAP*Taq* or pilot oligonucleotide as indicated in Figure 13B.

Strikingly, in the case of RNA cleavage, a 3' arm is not required for the pilot oligonucleotide. It is very unlikely that this cleavage is due to previously described RNaseH, which would be expected to cut the RNA in several places along the 30 base-pair long RNA-DNA duplex. The 5' nuclease of DNAP*Taq* is a structure-specific RNaseH that cleaves the RNA at a single site near the 5' end of the heteroduplexed region.

It is surprising that an oligonucleotide lacking a 3' arm is able to act as a pilot in directing efficient cleavage of an RNA target because such oligonucleotides are unable to direct efficient cleavage of DNA targets using native DNAPs. However, some 5' nucleases of the present invention (for example, clones E, F and G of Figure 4) can cleave DNA in the absence of a 3' arm. In other words, a non-extendable cleavage structure is not required for specific cleavage with some 5' nucleases of the present invention derived from thermostable DNA polymerases.

We tested whether cleavage of an RNA template by DNAP*Taq* in the presence of a fully complementary primer could help explain why DNAP*Taq* is unable to extend a DNA oligonucleotide on an RNA template, in a reaction resembling that of reverse transcriptase. Another thermophilic DNAP, DNAP*Tth*, is able to use RNA as a template, but only in the presence of Mn⁺⁺, so we predicted that this enzyme would not cleave RNA in the presence of this cation. Accordingly, we incubated an RNA molecule with an appropriate pilot oligonucleotide in the presence of DNAP*Taq* or DNAP*Tth*, in buffer containing either Mg⁺⁺ or Mn⁺⁺. As expected, both enzymes cleaved the RNA in the presence of Mg⁺⁺. However, DNAP*Taq*, but not DNAP*Tth*, degraded the RNA in the presence of Mn⁺⁺. We conclude that the 5' nuclease activities of many DNAPs may contribute to their inability to use RNA as templates.

EXAMPLE 2

Generation Of 5' Nucleases From Thermostable DNA Polymerases

Thermostable DNA polymerases were generated which have reduced synthetic activity, an activity that is an undesirable side-reaction during DNA cleavage in the detection assay of the invention, yet have maintained thermostable nuclease activity. The result is a thermostable polymerase which cleaves nucleic acids DNA with extreme specificity.

Type A DNA polymerases from eubacteria of the genus *Thermus* share extensive protein sequence identity (90% in the polymerization domain, using the Lipman-Pearson method in the DNA analysis software from DNASTar, WI) and behave similarly in both polymerization and nuclease assays. Therefore, we have used the genes for the DNA polymerase of *Thermus aquaticus* (DNAPTaq) and *Thermus flavus* (DNAPTfl) as representatives of this class. Polymerase genes from other eubacterial organisms, such as *Thermus thermophilus*, *Thermus sp.*, *Thermotoga maritima*, *Thermosipho africanus* and *Bacillus stearothermophilus* are equally suitable. The DNA polymerases from these thermophilic organisms are capable of surviving and performing at elevated temperatures, and can thus be used in reactions in which temperature is used as a selection against non-specific hybridization of nucleic acid strands.

The restriction sites used for deletion mutagenesis, described below, were chosen for convenience. Different sites situated with similar convenience are available in the *Thermus thermophilus* gene and can be used to make similar constructs with other Type A polymerase genes from related organisms.